

A New Simple Derivation of Schrödinger Uncertainty Relation

Gustavo Rigolin*

*Departamento de Física da Matéria Condensada,
Instituto de Física Gleb Wataghin, Universidade Estadual de Campinas,
C.P.6165, cep 13084-971, Campinas, São Paulo, Brazil*

We show how the Schrödinger Uncertainty Relation for a pair of observables can be deduced using the Cauchy-Schwarz inequality plus successive applications of the commutation relation involving the two observables. Our derivation differs from the original one in the sense that we do not need the expansion of the product of these two observables in a sum of symmetrical and anti-symmetrical operators.

I. INTRODUCTION

In 1930 Erwin Schrödinger presented^{1,2} a lower bound for the product of the dispersion of two non-commuting observables. This lower bound, from now on called Schrödinger Uncertainty Relation (SUR), is more general than the usual Heisenberg Uncertainty Relation (HUR) taught in all Quantum Mechanics courses. In fact, HUR is a particular case of SUR.

In this article we present an alternative derivation of Schrödinger's relation. Different from the original one, we do not make use of the expansion of the product of two observables in a sum of symmetrical and anti-symmetrical operators.

In Sec. II we review in a modern notation Schrödinger's derivation and in Sec. III we present the new one.

II. SCHRÖDINGER'S DERIVATION

Let X and P be our two non-commuting observables: $[X, P] = XP - PX \neq 0$. Now we define the following two states

$$|\psi\rangle = X|\chi\rangle, \quad (1)$$

$$|\phi\rangle = P|\chi\rangle. \quad (2)$$

For the moment we assume $\langle X \rangle = \langle \chi|X|\chi \rangle = 0$ and $\langle P \rangle = \langle \chi|P|\chi \rangle = 0$. These quantities are, respectively, the mean values of X and P for a system described by the normalized state $|\chi\rangle$. Applying the Cauchy-Schwarz inequality for the states (1) and (2),

$$\langle \psi|\psi \rangle \langle \phi|\phi \rangle \geq |\langle \psi|\phi \rangle|^2 = \langle \psi|\phi \rangle \langle \phi|\psi \rangle, \quad (3)$$

we get,

$$\langle X^2 \rangle \langle P^2 \rangle \geq \langle XP \rangle \langle PX \rangle. \quad (4)$$

Remembering that the dispersions of X and P are $\Delta X = (\langle X^2 \rangle - \langle X \rangle^2)^{1/2}$ and $\Delta P = (\langle P^2 \rangle - \langle P \rangle^2)^{1/2}$, Eq. (4) becomes

$$(\Delta X)^2 (\Delta P)^2 \geq \langle XP \rangle \langle PX \rangle. \quad (5)$$

Now we make use of Schrödinger's ingenuity and write XP as a sum of symmetrical (S) and anti-symmetrical

(A) operators:

$$XP = \frac{XP + PX}{2} + \frac{XP - PX}{2} = S + A, \quad (6)$$

$$PX = \frac{XP + PX}{2} - \frac{XP - PX}{2} = S - A. \quad (7)$$

Using Eqs. (6) and (7) in Eq. (5) we obtain

$$\begin{aligned} (\Delta X)^2 (\Delta P)^2 &\geq \langle S + A \rangle \langle S - A \rangle \\ &\geq \langle S \rangle^2 - \langle A \rangle^2 \\ &\geq \frac{\langle \{X, P\} \rangle^2}{4} - \frac{\langle [X, P] \rangle^2}{4}, \end{aligned} \quad (8)$$

where $\{X, P\} = XP + PX$ is the anti-commutator of X and P . Moreover, since X and P are Hermitian operators (observables) we know that $[X, P] = iC$, where C is Hermitian and $i = \sqrt{-1}$. Therefore, Eq. (8) can be written as,

$$(\Delta X)^2 (\Delta P)^2 \geq \frac{\langle \{X, P\} \rangle^2}{4} + \frac{|[X, P]|^2}{4}. \quad (9)$$

Finally, if we had $\langle X \rangle \neq 0$ and $\langle P \rangle \neq 0$ the proof can be carried out by making the following substitutions:

$$X \longrightarrow X - \langle X \rangle, \quad (10)$$

$$P \longrightarrow P - \langle P \rangle. \quad (11)$$

Repeating the previous procedure we arrive at the general form of SUR:

$$(\Delta X)^2 (\Delta P)^2 \geq \left(\frac{\langle \{X, P\} \rangle}{2} - \langle X \rangle \langle P \rangle \right)^2 + \frac{|[X, P]|^2}{4}. \quad (12)$$

III. AN ALTERNATIVE DERIVATION

We now present a new way of deriving Eqs. (9) and (12) without employing Eqs. (6) and (7), while maintaining the same simplicity of the previous derivation. The key idea behind the following deduction lies on the convenient use of the commutator of X and P .

Again, we begin with the simplest situation, i. e. $\langle X \rangle = \langle P \rangle = 0$ and, as before, our starting point is

Eq. (5). Remembering the definition of the commutator of X and P we can write XP and PX as

$$XP = PX + [X, P], \quad (13)$$

$$PX = XP - [X, P]. \quad (14)$$

Using Eq. (13) we can write Eq. (5) as

$$(\Delta X)^2(\Delta P)^2 \geq \langle PX \rangle^2 + \langle [X, P] \rangle \langle PX \rangle. \quad (15)$$

Now using Eq. (14) in Eq. (5) we get

$$(\Delta X)^2(\Delta P)^2 \geq \langle XP \rangle^2 - \langle [X, P] \rangle \langle XP \rangle. \quad (16)$$

Finally, using simultaneously Eqs. (13) and (14) in Eq. (5) we obtain,

$$(\Delta X)^2(\Delta P)^2 \geq \langle XP \rangle \langle PX \rangle + \langle [X, P] \rangle \langle XP \rangle - \langle [X, P] \rangle \langle PX \rangle - \langle [X, P] \rangle^2. \quad (17)$$

To finish the proof we add Eqs. (5), (15), (16), and (17):

$$\begin{aligned} (\Delta X)^2(\Delta P)^2 &\geq \frac{(\langle XP \rangle + \langle PX \rangle)^2}{4} - \frac{\langle [X, P] \rangle^2}{4} \\ &\geq \frac{\langle \{X, P\} \rangle^2}{4} + \frac{|\langle [X, P] \rangle|^2}{4}. \end{aligned} \quad (18)$$

If $\langle X \rangle \neq 0$ and $\langle P \rangle \neq 0$ the previous proof also works and we get SUR given by Eq. (12) if we repeat the previous procedure using Eqs. (10) and (11). $\square$

Acknowledgments

The author thanks Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) for funding this research.

* Electronic address: rigolin@ifi.unicamp.br

¹ E. Schrödinger, Proceedings of The Prussian Academy of Sciences **19**, 296 (1930). This text is written in german.

² A. Angelow and M. C. Batoni, Bulg. J. Phys. **26**, 193

(1999). Also available at e-print: quant-ph/9903100. This article is an english translation of Ref. 1.